

QUANTUM-CORRELATED MEASUREMENTS RELATED TO THE DETERMINATION OF γ/ϕ_3

J. LIBBY^{*1}

¹*Department of Physics, Indian Institute of Technology Madras,
Chennai 600028, Tamil Nadu, India
libby@iitm.ac.in*

(Dated: 13th December 2010)

Abstract

Measurements of D^0 meson strong-phase parameters in quantum-correlated $\psi(3770) \rightarrow D^0 \overline{D}^0$ decays by the CLEO collaboration are presented. These measurements play an important role in the determination of the unitarity triangle angle γ/ϕ_3 from B -meson decays. Measurements of the strong-phase parameters for $D^0 \rightarrow K^0 \pi^+ \pi^-$, $D^0 \rightarrow K^0 K^+ K^-$, $D^0 \rightarrow K^- \pi^+ \pi^0$, and $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ decays are described along with their impact on the determination of γ/ϕ_3 .

^{*} Proceedings of an invited talk at CHARM 2010, Beijing, China, Oct 21-24 2010, given on behalf of the CLEO Collaboration.

I. INTRODUCTION

One of the primary goals of flavor physics is to determine the angle γ/ϕ_3 of the $b-d$ CKM triangle [1]. Aside from being the least well known angle of the unitarity triangle (UT), it can be determined in tree-level processes that have negligible contributions from beyond the standard model physics, unlike most other constraints on the UT [2]. Therefore, any disagreement between the tree-level measurement of γ/ϕ_3 with predictions derived from other measurements is a signature of new physics.

The most promising decay to determine γ/ϕ_3 at tree level is $B^- \rightarrow \widetilde{D}^0 K^-$ where \widetilde{D}^0 is a D^0 or \overline{D}^0 decaying to the same final state [3].¹ The sensitivity to γ/ϕ_3 arises from the interference between the decay $B^- \rightarrow D^0 K^-$ and the color and CKM-suppressed decay $B^- \rightarrow \overline{D}^0 K^-$. The most precise measurements [4, 5] of γ/ϕ_3 come from decays where $\widetilde{D}^0 \rightarrow K_S^0 h^+ h^-$ [6, 7]. Here, h is π or K . Other promising \widetilde{D}^0 final states are $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$ [8, 9]. All these measurements depend on parameters related to the decay of the D^0 meson. Knowledge of the D -decay parameters *a priori* can greatly improve the determination of γ/ϕ_3 . These proceedings summarise the measurements [10, 11] of these parameters made by the CLEO collaboration and estimates their impact on the determination of γ/ϕ_3 .

II. MEASUREMENT OF THE STRONG-PHASE PARAMETERS OF $D^0 \rightarrow K^0 h^+ h^-$ DECAYS

The sensitivity to γ/ϕ_3 in $B^- \rightarrow \widetilde{D}^0(K_S^0 h^+ h^-)K^-$ comes from studying differences between the $\widetilde{D}^0 \rightarrow K_S^0 h^+ h^-$ Dalitz plot for both B^- and B^+ decays. Current measurements of γ/ϕ_3 require a model of the $\widetilde{D}^0 \rightarrow K_S^0 h^+ h^-$ Dalitz plot, which is derived from flavor-tagged samples of $D^0 \rightarrow K_S^0 h^+ h^-$. The assumptions used to determine the model introduce a systematic uncertainty on γ/ϕ_3 which is estimated to be between 3° and 9° [4, 5]. This is significantly less than the current statistical uncertainty but it will be a limiting factor in future measurements [12, 13]. Therefore, it is desirable to perform the measurement in a model-independent manner. Such a method was proposed in Ref. [6] and has been developed significantly by Bondar and Poluektov [14]. The method requires determining yields in bins of the $\widetilde{D}^0 \rightarrow K_S^0 h^+ h^-$ Dalitz plot for B^- and B^+ decay, which depend on the B -decay parameters and two new parameters c_i and s_i , which are the amplitude-weighted averages over the bin of the cosine and sine of the difference in strong-phase difference, $\Delta\delta_D$, between Dalitz-plot points (m_-^2, m_+^2) and (m_+^2, m_-^2) . Here m_\pm is the invariant-mass of the $K_S^0 h^\pm$ pair. It can be shown [10, 14] that between 80% to 90% of the statistical sensitivity to γ/ϕ_3 of the unbinned method can be obtained by choosing bins corresponding to equal intervals of $\Delta\delta_D$ according to an amplitude model. An example of such a binning is shown in Fig. 1.

The values of c_i and s_i can be measured in quantum-correlated $D^0 \overline{D}^0$ decays of the $\psi(3770)$. The $D^0 \overline{D}^0$ are produced in a $C = -1$ state. Therefore, if one D meson decays to a CP -eigenstate the other D -meson is in the opposite CP -eigenstate. The difference between CP -even and CP -odd tagged Dalitz plots in each bin is related to the c_i parameters. In addition, the Dalitz plot of quantum-correlated events where both D -mesons decay to $K_S^0 h^+ h^-$ is sensitive to both c_i and s_i . The strong-phase parameters for the decay $D^0 \rightarrow$

¹ Here and throughout this paper the charge-conjugate state is implied unless otherwise stated.

$K_L^0 h^+ h^-$ (c'_i and s'_i) are closely related to c_i and s_i such that using decays of the type $K_S^0 h^+ h^-$ *vs.* $K_L^0 h^+ h^-$ greatly improve the precision on c_i and s_i .

The CLEO-c experiment [15] collected $e^+e^- \rightarrow \psi(3770) \rightarrow D\bar{D}$ data corresponding to an integrated luminosity of 818 pb^{-1} . The fact that all particles arise from D -meson decay in the final state leads to both D mesons being reconstructed exclusively with high efficiency and purity. For $D^0 \rightarrow K^0 \pi^+ \pi^-$ ($D^0 \rightarrow K^0 K^+ K^-$) decay the numbers of CP -tagged and $K^0 h^+ h^-$ *vs.* $K^0 h^+ h^-$ candidates selected are 1661 and 1674 (219 and 335), respectively.

A maximum-likelihood fit is performed to the bin yields of the CP -tagged and $K^0 h^+ h^-$ *vs.* $K^0 h^+ h^-$ events to extract $c_i^{(\prime)}$ and $s_i^{(\prime)}$. The results are presented in detail elsewhere [10]. The values of $c_i^{(\prime)}$ and $s_i^{(\prime)}$ are determined for several binning variations for both $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $D^0 \rightarrow K_S^0 K^+ K^-$. These binnings allow flexibility given different scenarios for the amount of B data and the background environment. The measured values of c_i and s_i are found to be in reasonable agreement with the values predicted by the amplitude models presented in Refs. [4, 16]. The largest systematic uncertainties arise from the modelling of the background. However, none of these measurements are systematically limited.




FIG. 1: Dalitz-plot binning for $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ in region of similar $\Delta\delta_D$.

III. MEASUREMENT OF THE COHERENCE FACTOR AND STRONG PHASE DIFFERENCES IN $D^0 \rightarrow K^- \pi^+ \pi^0$ AND $D^0 \rightarrow K^- \pi^+ \pi^-$

The rate of decays $B^- \rightarrow \widetilde{D}^0(K^+ \pi^-)K^-$ is particularly sensitive to γ/ϕ_3 because the two interfering amplitudes are of similar size due to the doubly-Cabibbo suppressed (DCS) D^0 decay coming from the favored B^- amplitude [8]. The rate depends not only on γ/ϕ_3 but on the strong-phase difference between the Cabibbo-favored and DCS $\widetilde{D}^0 \rightarrow K^+ \pi^-$ decays. The measurement of this parameter by the CLEO collaboration is described elsewhere in these proceedings [18].

Via the same mechanism there is potential sensitivity to γ/ϕ_3 from $B^- \rightarrow \widetilde{D}^0 K^-$, where $\widetilde{D}^0 \rightarrow K^+ \pi^- \pi^0$ or $\widetilde{D}^0 \rightarrow K^+ \pi^- \pi^- \pi^+$ [9]. These modes have significantly larger branching fractions than $\widetilde{D}^0 \rightarrow K^+ \pi^-$ [17]. However, the dynamics are more complicated because there is variation of the strong-phase difference over the multi-body phase-space. This leads to the introduction of a new parameter referred to as the coherence factor R_F ($F = K\pi\pi^0$ or $K3\pi$), which multiplies the interference term sensitive to γ/ϕ_3 . The value of R_F can vary between zero and one. If there is only a single intermediate resonance or a few non-interfering resonances the coherence factor will be close to one and the decay will behave just like $\widetilde{D}^0 \rightarrow K^+ \pi^-$. If there are many overlapping intermediate resonances the coherence factor will tend toward zero, limiting the sensitivity to γ/ϕ_3 . However, even if there is limited sensitivity to the phases when $R \sim 0$ there is enhanced sensitivity to the magnitude of the amplitude ratio between the $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow \overline{D}^0 K^-$ decays; improved knowledge of this parameter will then lead to better overall sensitivity to γ/ϕ_3 in a global fit to all $B^- \rightarrow \widetilde{D}^0 K^-$ decays.

The values of R_F and the average-strong phase difference δ_D^F have been measured by CLEO-c [11]. Sensitivity comes from the quantum-correlated $D^0 \overline{D}^0$ events with F tagged by either CP -eigenstates or $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$, where the tag kaon charge is the same as the signal. A χ^2 fit to the yields gives: $R_{K\pi\pi^0} = 0.84 \pm 0.07$, $\delta_D^{K\pi\pi^0} = (227^{+14}_{-17})^\circ$, $R_{K3\pi} = 0.33^{+0.26}_{-0.23}$, and $\delta_D^{K3\pi} = (114^{+26}_{-23})^\circ$. Figure 2 shows the 1σ , 2σ , and 3σ regions of $(R_{K\pi\pi^0}, \delta_D^{K\pi\pi^0})$ parameter space; the coherence of $D^0 \rightarrow K^- \pi^+ \pi^0$ is clearly observed. The impact of these results on the measurements of γ/ϕ_3 is discussed in the following section.

IV. IMPACT OF RESULTS ON THE MEASUREMENT OF γ/ϕ_3

The determination of the c_i and s_i in quantum-correlated D -decay allows the measurement of γ/ϕ_3 without a model induced systematic uncertainty. However, this is replaced by uncertainty due to the limited statistics used to measure c_i and s_i at CLEO-c. This uncertainty is estimated to be between 1.7° and 3.9° (3.2° and 3.9°) depending on the binning of the $D^0 \rightarrow K_S^0 \pi^+ \pi^+$ ($D^0 \rightarrow K_S^0 K^+ K^-$) Dalitz plot. The systematic uncertainty on γ/ϕ_3 can be reduced by about a factor of three if BES-III collects 10 fb^{-1} of integrated luminosity at the $\psi(3770)$ resonance [19]. This would reduce the uncertainty to the order of 1° , which would not only be adequate for LHCb but also for future higher luminosity facilities [13, 20].

The impact of the measurements of R_F and δ_D^F at LHCb is evaluated using the yield estimates for $B^- \rightarrow \widetilde{D}^0(K^\pm \pi^\mp)K^-$ and $B^- \rightarrow \widetilde{D}^0(K^\pm \pi^\mp \pi^+ \pi^-)K^-$ decays in a dataset corresponding to 2 fb^{-1} of integrated luminosity at a center-of-mass energy of 14 TeV [21]. In addition, the yield of $B^- \rightarrow \widetilde{D}^0(K^\pm \pi^\mp \pi^0)$ is assumed to be half that of $B^- \rightarrow \widetilde{D}^0(K^\pm \pi^\mp \pi^+ \pi^-)K^-$ with the same level of background reflecting the difficulties

associated with π^0 reconstruction in the hadronic environment. The sensitivity to γ/ϕ_3 from LHCb data alone is 9.7° . Including the CLEO-c constraints on R_F and δ_D^F this improves to 7.5° . The introduction of the CLEO-c constraints is equivalent to 70% more LHCb data. This clearly illustrates the power of quantum-correlated measurements in aiding the determination of γ/ϕ_3 . BES-III data could lead to at least a further 10% reduction of the uncertainty on γ/ϕ_3 .

In conclusion, the first quantum-correlated measurements of strong-phase parameters of D -decay at CLEO-c have been presented and their positive impact on the determination of γ/ϕ_3 at LHCb has been illustrated. Further improvements are possible by exploiting the larger sample of quantum-correlated decays that will be available at BES-III. Furthermore, there are other \bar{D}^0 decay modes of interest to the measurement of γ/ϕ_3 for which the strong-phase parameters have yet to be determined: $K_S^0\pi^+\pi^-\pi^0$, $\pi^+\pi^-\pi^0$, $K_S^0K^\pm\pi^\mp$, and $K^+K^-\pi^+\pi^-$.




FIG. 2: The 1σ , 2σ , and 3σ allowed regions of $(R_{K\pi\pi^0}, \delta_D^{K\pi\pi^0})$ parameter space.

Acknowledgments

I would like to thank the CHARM 2010 organisers for their financial assistance that allowed my participation in an excellent conference.

-
- [1] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
 - [2] CKMfitter Group, (A. Höcker *et al.*), *Eur. Phys. J. C* **21**, 225 (2001); CKMfitter Group (J. Charles *et al.*), *Eur. Phys. J. C* **41**, 1 (2005), and updates at <http://ckmfitter.in2p3.fr/>; M. Ciuchini *et al.*, *J. High Energy Phys.* 0107 (2001) 013; UTFit Collaboration (M. Bona *et al.*), *J. High Energy Phys.* 081 (2006) 10, and updates at <http://www.utfit.org/>.
 - [3] M. Gronau and D. Wyler, *Phys. Lett. B* **265**, 172 (1991); M. Gronau and D. London, *Phys. Lett. B* **253**, 483 (1991).
 - [4] Belle Collaboration (A. Poluektov *et al.*), *Phys. Rev. D* **81**, 112002 (2010).
 - [5] BABAR Collaboration, (P. del Amo Sanchez *et al.*), *Phys. Rev. Lett.* **105**, 121801 (2010).
 - [6] A. Giri, Y. Grossman, A. Soffer, and J. Zupan, *Phys. Rev. D* **68**, 054018 (2003).
 - [7] A. Bondar, *Proceedings of BINP Special Analysis Meeting on Dalitz Analysis*, 24-26 Sep. 2002, unpublished.
 - [8] D. Atwood, I. Dunietz, and A. Soni, *Phys. Rev. Lett.* **78**, 3257 (1997); D. Atwood, I. Dunietz, and A. Soni, *Phys. Rev. D* **63**, 036005 (2001).
 - [9] D. Atwood and A. Soni, *Phys. Rev. D* **68**, 033003 (2003).
 - [10] CLEO Collaboration (J. Libby *et al.*), arXiv:1010.2817 [hep-ex], accepted by *Phys. Rev. D*.
 - [11] CLEO Collaboration (N. Lowrey *et al.*), *Phys. Rev. D* **80**, 031105 (2009).
 - [12] J. Libby, CERN-LHCb-2007-141; V. Gibson, C. Lazzeroni, and Y.-Y. Li, CERN-LHCb-2008-028.
 - [13] T. Aushev *et al.*, KEK-Report 2009-12; M. Bona *et al.*, SLAC-R-856.
 - [14] A. Bondar and A. Poluektov, *Eur. Phys. J. C* **47**, 347 (2006); A. Bondar and A. Poluektov, *Eur. Phys. J. C* **55**, 51 (2008).
 - [15] CLEO Collaboration (Y. Kubota *et al.*), *Nucl. Instrum. Methods Phys. Res., Sect. A* **320**, 66 (1992); D. Peterson *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **478**, 142 (2002); M. Artuso *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **554**, 147 (2005).
 - [16] BABAR Collaboration (B. Aubert *et al.*), *Phys. Rev. D* **78**, 034023 (2008).
 - [17] Particle Data Group (K. Nakamura *et al.*), *J. Phys. G* **37**, 075021 (2010).
 - [18] D. Asner, these proceedings.
 - [19] H. Liu, these proceedings.
 - [20] LHCb Collaboration, CERN-LHCb-2008-019.
 - [21] LHCb Collaboration (B. Adeva *et al.*), LHCb-PUB-2009-029.